



Advancing Sea Level Change Science with SWOT

R. Steven Nerem and Dallas Masters, University of Colorado



Global mean sea level change variations over the last 20 years

Regional sea level trends over the last 20 years



Measuring Sea Level Change

- SWOT can potentially continue and augment the GMSL climate record, densify the regional sea level record, and pioneer the coastal sea level record.
- The nominal SWOT inclination (78°) will provide coverage of the polar regions not currently covered by the reference altimeter missions ($\pm 66^\circ$ - TOPEX, Jason-1, Jason-2).
- Regional sea level changes measured by SWOT in the polar regions can provide improved observability of the sea level fingerprints associated with ice sheet mass loss.
- SWOT has the potential to provide improved measurements of coastal sea level, which is critical for assessing the socio-economic impacts of sea level change.
- SWOT will help us understand the relation between sea level and terrestrial water cycle.

Our Contribution to the SWOT SDT

- We will work with the SDT to understand how SWOT might contribute to sea level change science.
- We will investigate how the SWOT wide-swath measurements might improve estimates of regional and coastal sea level change.
- We will develop approaches to calibrate and validate the SWOT nadir and wide-swath sea level measurements using comparisons to other altimeters, tide gauge data, etc.
- We will assess the ability of the tide gauge network for monitoring instrument drifts in the SWOT measurements.
- We will investigate other novel approaches to calibrate/validate the wide-swath sea level measurements.
- We can assist with the design of the SWOT orbit as needed.

Phase A – SWOT Sea Level Issues

- For sea level science from the nadir altimeter (and for understanding cross-calibrations with other altimeter missions), it would still be desirable to have SWOT fly a ± 1 km repeat orbit.
- For sea level science, we believe high-resolution data is desired for sea level studies in coastal regions, but probably not needed in the open ocean for sea level change studies.
- A high-level gridded sea level product would be desirable, including a gridded mean sea surface that incorporates the SWOT wide-swath data.
- The footprint of the radiometer and correctly assigning land flags will be crucial for coastal SSH measurements.
- Improved coastal tide and mean sea surface models will be essential.
- Validate with coastal tide gauges, AirSWOT, etc.